1L HER2+ GEA

Reimagine tomorrow

with TEVIMBRA + zanidatamab-hrii + chemotherapy

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

In a clinical trial, people with unresectable locally advanced or metastatic HER2+ GEA lived longer with TEVIMBRA + zanidatamab-hrii + chemotherapy than with trastuzumab + chemotherapy

- 26-month median overall survival* for the 302 people taking TEVIMBRA + zanidatamab-hrii + chemotherapy (vs 19 months for the 308 people taking trastuzumab + chemotherapy)

*Median overall survival is the length of time in a clinical trial from when treatment starts to when half of those enrolled in a treatment combination are still alive.

HER2, human epidermal growth factor receptor 2; GEA, gastroesophageal adenocarcinoma.

What is TEVIMBRA?

TEVIMBRA® (tislelizumab-jsgr) is a prescription medicine used to treat adults with:

- cancer of the stomach (gastric cancer), or cancer where the esophagus joins the stomach (gastroesophageal junction cancer), or of the tube that connects your throat to your stomach (esophageal cancer).
  - TEVIMBRA may be used in combination with zanidatamab-hrii and fluoropyrimidine- and platinum-containing chemotherapy as your first treatment when your gastric, gastroesophageal, or esophageal adenocarcinoma:
    - cannot be removed with surgery or has spread to other parts of the body (metastatic), **and**
    - is human epidermal growth factor receptor 2 (HER2)-positive (IHC 3+ or IHC 2+/ISH+).

    You should also refer to the Medication Guide that comes with zanidatamab-hrii.

It is not known if TEVIMBRA is safe and effective in children.

IMPORTANT SAFETY INFORMATION

TEVIMBRA works with your immune system. It can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended.

Please see additional Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.



GET STARTED

Table of contents

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

Click on the buttons below to learn more about a topic:

UNDERSTANDING THE DISEASE AND TREATMENT

3 What is HER2+ GEA?

4 About TEVIMBRA + zanidatamab-hrii + chemotherapy

5 Results with TEVIMBRA + zanidatamab-hrii + chemotherapy

GETTING STARTED WITH TREATMENT

7 What is an IV infusion?

8 Preparing for your first appointment

9 What to bring to infusions

10 What to expect during infusions

TRACKING HOW YOU'RE FEELING

11 Managing side effects

14 Managing diet

HELPFUL SUPPORT AND RESOURCES

15 Cancer community resources

16 myBeOne Patient Support

IMPORTANT SAFETY INFORMATION

17 Important Safety Information

The information provided is meant for informational purposes only. It is not meant to replace your physician's medical advice.

Please see Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

What is unresectable locally advanced or metastatic HER2+ GEA?

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

GEA stands for gastroesophageal adenocarcinoma. It starts when healthy cells within the digestive organs become cancerous and begin to grow and spread. GEA includes:

- Gastric cancer (cancer of the stomach)
- Esophageal cancer (cancer of the tube that connects the throat to the stomach)
- Gastroesophageal junction cancer (cancer where the esophagus joins the stomach)

Being HER2-positive (HER2+) means that those cancer cells are making too much of a protein called HER2. This can cause the cancer to grow and spread more quickly.

Unresectable locally advanced GEA means that the cancer has spread to nearby areas of the body and cannot be safely removed with surgery. Having metastatic GEA means that the cancer has spread to other parts of the body.

Actor portrayals.

Please see Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

TEVIMBRA + zanidatamab + chemotherapy are thought to work together to attack cancer from multiple angles

TEVIMBRA® (tislelizumab-jsgr) Injection, for intravenous use 10 mg/mL

Each medicine in this combination works in a distinct way to help attack cancer cells:

TEVIMBRA

Helps your immune system turn your T cells back on

+

Chemotherapy

Works to destroy cancer cells throughout your body

+

Zanidatamab

Binds to HER2 proteins on cancer cells to stop their growth and help your immune system attack them

T cells are a type of white blood cell in your immune system. Their job is to find and destroy disease and infection throughout your body. But when you have cancer, your T cells have been "switched off." This allows the cancer to grow and spread more quickly.

IMPORTANT SAFETY INFORMATION (cont'd)

What signs and symptoms of immune system problems can happen with TEVIMBRA?

Contact your healthcare provider right away if you develop any new or worsening symptoms including:

- Lung problems: New or worsening cough, shortness of breath, and chest pain.
- Intestinal problems: Diarrhea or more bowel movements than usual; stools that are black, tarry, sticky, or have blood or mucus; and severe stomach-area pain or tenderness.

Please see additional Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

In people with unresectable locally advanced or metastatic HER2+ GEA,

TEVIMBRA + zanidatamab-hrii + chemotherapy may help people live longer than trastuzumab + chemotherapy

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

TEVIMBRA + zanidatamab-hrii + chemotherapy was studied in a clinical trial of 914 people with advanced HER2+ GEA. People enrolled received one of three treatment combinations, two of which are compared below.

With TEVIMBRA + zanidatamab-hrii + chemotherapy, people typically lived longer than with trastuzumab + chemotherapy

AT ABOUT 26 MONTHS, HALF OF THE 302 PEOPLE receiving **TEVIMBRA** + zanidatamab-hrii + chemotherapy **WERE STILL ALIVE**

VS

AT ABOUT 19 MONTHS, HALF OF THE 308 PEOPLE receiving trastuzumab + chemotherapy **WERE STILL ALIVE**

This data is based on median overall survival, which is the length of time in a clinical trial from when treatment starts to when half of those enrolled in a treatment combination are still alive.

IMPORTANT SAFETY INFORMATION (cont'd)

What signs and symptoms of immune system problems can happen with TEVIMBRA? (cont'd)

- Liver problems: Yellowing of your skin or the whites of your eyes; severe nausea or vomiting; pain on the right side of your stomach area; dark urine; and bleeding or bruising more easily than normal.
- Hormone gland problems: Headaches that will not go away or unusual headaches; eye sensitivity to light; eye problems; rapid heartbeat; increased sweating; extreme tiredness; weight gain or weight loss; feeling more hungry or thirsty than usual; urinating more often than usual; hair loss; feeling cold; constipation; voice gets deeper; dizziness or fainting; and changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness.

Please see additional Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

TEVIMBRA + zanidatamab-hrii + chemotherapy helped shrink tumors and lowered the risk of cancer progression and death*

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

In the same clinical trial:

28% lower risk

People taking TEVIMBRA + zanidatamab-hrii + chemotherapy had a **28% lower relative risk of death** compared to those taking trastuzumab + chemotherapy

2X as likely

At 2 years, people taking TEVIMBRA + zanidatamab-hrii + chemotherapy were **more than twice as likely to be alive** without their cancer getting worse compared to people taking trastuzumab + chemotherapy (38% vs 16%)

67% of people

67% of the 302 people taking TEVIMBRA + zanidatamab-hrii + chemotherapy responded to treatment—this means that **their tumors shrank or even disappeared completely** (vs 63% of the 308 people talking trastuzumab + chemotherapy)

- Half of those who responded to TEVIMBRA + zanidatamab-hrii + chemotherapy **did so for 19 months or more** (compared to 8 months for those taking trastuzumab + chemotherapy)

***Compared to trastuzumab + chemotherapy.**

IMPORTANT SAFETY INFORMATION (cont'd)

What signs and symptoms of immune system problems can happen with TEVIMBRA? (cont'd)

- Kidney problems: Decrease in your amount of urine, blood in your urine, swelling in your ankles, and loss of appetite.
- Skin problems: Rash; itching; skin blistering or peeling; painful sores or ulcers in your mouth or nose, throat, or genital area; fever or flu-like symptoms; and swollen lymph nodes.

Please see additional Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

What is an IV infusion?

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

An intravenous (or IV) infusion is a way to deliver medicine slowly into your bloodstream through a vein. TEVIMBRA and zanidatamab-hrii are given by IV infusion, while chemotherapy may be given by IV infusion, oral pills, or a combination of both.

Infusions are a very common way to deliver medication, especially cancer medication. Your infusions may be given at your doctor's office, an infusion center, or another healthcare location.

IMPORTANT SAFETY INFORMATION (cont'd)

What signs and symptoms of immune system problems can happen with TEVIMBRA? (cont'd)

Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with TEVIMBRA. Contact your healthcare provider right away if you have any new or worsening signs or symptoms which may include chest pain, irregular heartbeat, shortness of breath, swelling of ankles; confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs; double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight; persistent or severe muscle pain or weakness, muscle cramps; low red blood cells, bruising.

Please see additional Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

Preparing for your first appointment

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

Taking a few steps before your infusion appointment can help you feel ready to begin treatment.

Confirm

Confirm your appointment with your healthcare provider. Be sure to notify your healthcare provider if you have any medical conditions or if you are pregnant or breastfeeding. Also tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Ask

Here are some questions you may want to ask your healthcare provider.

Before the appointment:

- Are there any foods or drinks I should avoid?
- Should I take my medications in the morning before treatment?
- What will my treatment schedule look like?

At the appointment:

- How long does the infusion take?
- Can a friend or family member stay with me?

After the appointment:

- What are common side effects of the treatment? Who should I call if I have side effects?
- Will I be able to drive myself to future appointments?

Plan

Make sure to arrive early for your first appointment to allow time for paperwork. For your infusion, be sure to wear comfortable clothes, including a button-down shirt with sleeves you can roll up past your elbow.

For caregivers: You can help your loved one prepare for infusion by:

- **Offering to provide or schedule transportation** to and from the infusion center
- **Coordinating with the healthcare team** to keep up to date with the treatment schedule
- **Keeping a notebook** for recording information about treatment, symptoms, and medications

Please see Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

What to bring to your infusion appointment

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

Being prepared and comfortable during your infusion appointment is important. Here is a list of items that may help you:

- An insurance card and any necessary paperwork, such as a list of current medications
- A sweater, in case the infusion center is chilly
- A pillow and blanket
- A book or tablet to help pass the time
- Headphones or earplugs
- Healthy snacks or drinks

Be sure to ask your healthcare team if you have any questions about what items you are allowed to bring with you.

Actor portrayal.

Please see Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

What to expect during your infusions

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

Your first infusion

- **When you arrive, a healthcare provider will get you settled** in the infusion area, set up your IV line, and start your treatment
- **Your infusion appointment will likely take several hours.** This is to allow time for premedications, hydration, and to ensure that each individual medicine (TEVIMBRA, zanidatamab-hrii, and chemotherapy) can be administered safely. The healthcare provider will continue to check on you to make sure you're feeling okay and are still comfortable. During this time you can read a book, listen to music or a podcast, or simply relax
- **Make sure to tell your healthcare provider if you become uncomfortable** during the infusion so they can offer help and monitor your needs. If you develop certain side effects, your doctor may pause, delay, or stop your treatment
- **When your first infusion ends, you may need to stay in the office** an extra 30 minutes for observation with your healthcare team. This helps ensure that you are tolerating the treatment well

Your next infusions

- **Your future infusions will be similar to your** first and will follow a regular 2, 3, 4 or 6-week schedule, depending on the dose you're receiving. This schedule will be determined by your doctor
- **You will continue treatment as long as it is working** and your body is handling it well
- **If you miss an appointment,** call your healthcare provider right away to reschedule

For caregivers: Support and companionship go a long way during the infusion sessions. If your schedule allows, consider keeping your loved one company while they receive their infusion.

Diarrhea is a possible side effect of this treatment combination. To help manage this side effect, your doctor may prescribe a common anti-diarrheal medication during your first treatment cycle, which some people may need to continue taking. Your doctor will determine what's best based on your personal experience.

Be sure to reach out to your healthcare team if you have any questions or concerns.

IMPORTANT SAFETY INFORMATION (cont'd)

What problems can occur during infusion?

Infusion reactions can sometimes be severe or life-threatening. Signs or symptoms of infusion reactions may include chills or shaking, itching or rash, flushing, shortness of breath or wheezing, dizziness, feeling like passing out, fever, or back or neck pain.

Please see additional Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

It's important to watch for side effects

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

You know your body best.

Paying attention to how you feel helps your healthcare team recognize and manage side effects early.

Be sure to:

- **Communicate any side effects to your healthcare provider**
- **Write down any side effects you feel in a treatment journal**
- **Talk to your friends and family about side effects**
- **Reach out to support groups**

The most common side effects associated with TEVIMBRA + zanidatamab-hrii + chemotherapy are:

- diarrhea
- nausea
- decreased red blood cell count (anemia)
- decreased appetite
- vomiting
- decreased potassium in your blood
- tiredness
- rash
- decreased white blood cell count
- decreased platelet count
- numbness, pain, tingling, or burning in your hands or feet
- infusion reactions
- increase in certain liver blood tests

See the following pages for more possible side effects and talk to your doctor should they occur.

Please see Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

Possible side effects that may occur

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

Call your doctor if you experience any of the following:

Lung problems
- New or worsening cough
- Shortness of breath
- Chest pain

Intestinal problems
- Diarrhea (loose stools) or more bowel movements than usual
- Stools that are black, tarry, sticky, or have blood or mucus
- Severe stomach-area (abdomen) pain or tenderness

Liver problems
- Yellowing of your skin or the whites of your eyes
- Severe nausea or vomiting
- Pain on the right side of your stomach area (abdomen)
- Dark urine (tea colored)
- Bleeding or bruising more easily than normal

Hormone gland problems
- Headaches that will not go away or unusual headaches
- Eye sensitivity to light
- Eye problems
- Rapid heartbeat
- Increased sweating
- Extreme tiredness
- Weight gain or weight loss
- Feeling more hungry or thirsty than usual
- Urinating more often than usual
- Hair loss
- Feeling cold
- Constipation
- Your voice gets deeper
- Dizziness or fainting
- Changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness

Kidney problems
- Decrease in your amount of urine
- Blood in your urine
- Swelling in your ankles
- Loss of appetite

Skin problems
- Rash
- Itching
- Skin blistering or peeling
- Painful sores or ulcers in your mouth or in your nose, throat, or genital area
- Fever or flu-like symptoms
- Swollen lymph nodes

For caregivers: You can help observe and take note of any side effects your loved one may be experiencing. Encourage them to be honest about their symptoms. Proactively talk with your loved one's healthcare team about any potential side effects.

Please see Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

Continued on next page →

Possible side effects that may occur (cont'd)

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

Call your doctor if you experience any of the following:



Immune system problems in other organs

- Chest pain, irregular heartbeat, shortness of breath, swelling of ankles
- Confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight
- Persistent or severe muscle pain or weakness, muscle cramps
- Low red blood cells, bruising

Reactions during or shortly after the infusion that can sometimes be severe or life-threatening

- Chills or shaking
- Itching or rash
- Flushing
- Shortness of breath or wheezing
- Dizziness
- Feeling like passing out
- Fever
- Back or neck pain

IMPORTANT SAFETY INFORMATION (cont'd)

What if I have a transplanted organ or tissue; or bone marrow transplant?

Rejection of a transplanted organ or tissue. Your healthcare provider should tell you what signs and symptoms you should report and monitor you, depending on the type of organ or tissue transplant that you have had. There may be complications, including graft-versus-host-disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells. These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with TEVIMBRA. Your healthcare provider will monitor you for these complications.

Getting medical treatment right away may help keep these problems from becoming more serious. Your healthcare provider will check you for these problems during your treatment with TEVIMBRA. Your healthcare provider may treat you with corticosteroid or hormone replacement medicines, or may delay or completely stop treatment with TEVIMBRA if you have severe side effects.

Please see additional Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

Managing your diet during treatment

TEVIMBRA® (tislelizumab-jsgr) Injection, for intravenous use 10 mg/mL

HER2+ GEA often impacts your appetite, digestion, or ability to swallow. Your care team may suggest eating smaller, more frequent meals and choosing high-protein, easy-to-digest foods, such as:

- Oatmeal
- Pureed soups
- Mashed potatoes
- Scrambled eggs
- Yogurt and pudding
- Soft rice and pasta
- Smoothies

For caregivers: Ask your loved one's healthcare team or dietitian for meal ideas that fit their specific dietary needs.

IMPORTANT SAFETY INFORMATION (cont'd)

Before receiving TEVIMBRA, tell your healthcare provider about all of your medical conditions, including if you have immune system problems; received an organ or tissue transplant; received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic); received radiation treatment to your chest area; or a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome; or are pregnant, plan to become pregnant, or are breastfeeding or plan to breastfeed. Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see additional Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

Cancer community resources

TEVIMBRA® (tislelizumab-jsgr) Injection, for intravenous use 10 mg/mL

You and your caregivers are not alone. Organizations are here to help.

Visit the websites below for information about transportation, education, counseling services, and more:

HOPE FOR STOMACH CANCER	Hope For Stomach Cancer is a 501(c)(3) nonprofit organization that provides resources to patients, caregivers, and loved ones while promoting early detection and prevention to the general and medical communities. The organization creates and facilitates programs that enable those affected by stomach cancer to take actionable steps to live with each phase of the disease. Their vision is to bridge the gap between research and patient care.	**stocan.org**
GASTRIC CANCER FOUNDATION	The Gastric Cancer Foundation is a national nonprofit organization dedicated to improving the lives of people affected by gastric (stomach) cancer and working with leading researchers on treatments. The foundation website offers a comprehensive range of resources for those impacted by the disease. Online offerings include a confidential clinical trial finder, a secure community forum, and a video series with nutritional tips and recipes tailored to the unique needs of this population, among other valuable tools.	**gastriccancer.org**
CANCERcare	Cancer*Care* is the leading national organization providing free professional support services and information to help manage the emotional, practical, and financial challenges of cancer. Resources include counseling, educational workshops, and copayment assistance provided by social workers and world-leading cancer experts.	**cancercare.org** **800-813-4673**
CANCER SUPPORT COMMUNITY	CSC is a global nonprofit network with 190 locations, including CSC and Gilda's Club centers as well as healthcare partnerships. These locations, along with a toll-free helpline, digital services, and award-winning education materials, provide $50 million in free support services to patients and families each year.	**cancersupportcommunity.org** **272-867-5309**
American Cancer Society	The American Cancer Society (ACS) is a nationwide, community-based volunteer health organization that aims to eliminate cancer as a major health problem. ACS provides information and support, such as transportation and lodging, to people in every community who have been touched by cancer.	**cancer.org** **800-227-2345**

BeOne Medicines does not control or endorse third-party organizations. As such, BeOne Medicines makes no representation as to the accuracy or any other aspect of the information supplied by these organizations or contained on these websites.

Please see Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

The myBeOne Support program is here for you throughout your treatment journey

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

myBeOneSupport® can pair you with a dedicated Oncology Nurse Advocate. They can provide personalized support for eligible patients and caregivers throughout treatment with TEVIMBRA.

Additional services include:

- **Simplifying access**
- **Educating patients and caregivers about treatment and disease**
- **Connecting patients to independent organizations* that may be able to assist with day-to-day living support**

*Independent, third-party organizations have their own rules for eligibility. BeOne Medicines has no involvement in their decision-making or eligibility criteria. This information is nonexhaustive and not meant as an endorsement from BeOne Medicines.

myBeOneSupport®

For any questions, or to help you get started, call 1-833-234-4363, Monday through Friday, 8 AM–8 PM ET, to speak with an Oncology Nurse Advocate. You can also visit myBeOneSupport.com.

myBeOneSupport® is not insurance and does not guarantee product coverage or reimbursement.

Please see Important Safety Information on pages 17-18 and Medication Guide for TEVIMBRA.

IMPORTANT SAFETY INFORMATION

TEVIMBRA® (tislelizumab-jsgr) Injection, for intravenous use 10 mg/mL

TEVIMBRA works with your immune system. It can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended.

What signs and symptoms of immune system problems can happen with TEVIMBRA?

Contact your healthcare provider right away if you develop any new or worsening symptoms including:

- Lung problems: New or worsening cough, shortness of breath, and chest pain.
- Intestinal problems: Diarrhea or more bowel movements than usual; stools that are black, tarry, sticky, or have blood or mucus; and severe stomach-area pain or tenderness.
- Liver problems: Yellowing of your skin or the whites of your eyes; severe nausea or vomiting; pain on the right side of your stomach area; dark urine; and bleeding or bruising more easily than normal.
- Hormone gland problems: Headaches that will not go away or unusual headaches; eye sensitivity to light; eye problems; rapid heartbeat; increased sweating; extreme tiredness; weight gain or weight loss; feeling more hungry or thirsty than usual; urinating more often than usual; hair loss; feeling cold; constipation; voice gets deeper; dizziness or fainting; and changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness.
- Kidney problems: Decrease in your amount of urine, blood in your urine, swelling in your ankles, and loss of appetite.
- Skin problems: Rash; itching; skin blistering or peeling; painful sores or ulcers in your mouth or nose, throat, or genital area; fever or flu-like symptoms; and swollen lymph nodes.

Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with TEVIMBRA. Contact your healthcare provider right away if you have any new or worsening signs or symptoms which may include chest pain, irregular heartbeat, shortness of breath, swelling of ankles; confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs; double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight; persistent or severe muscle pain or weakness, muscle cramps; low red blood cells, bruising.

What problems can occur during infusion?

Infusion reactions can sometimes be severe or life-threatening. Signs or symptoms of infusion reactions may include chills or shaking, itching or rash, flushing, shortness of breath or wheezing, dizziness, feeling like passing out, fever, or back or neck pain.

Please see additional Important Safety Information on the following page and Medication Guide for TEVIMBRA.

IMPORTANT SAFETY INFORMATION (cont'd)

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

What if I have a transplanted organ or tissue; or bone marrow transplant?

Rejection of a transplanted organ or tissue. Your healthcare provider should tell you what signs and symptoms you should report and monitor you, depending on the type of organ or tissue transplant that you have had. There may be complications, including graft-versus-host-disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells. These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with TEVIMBRA. Your healthcare provider will monitor you for these complications.

Getting medical treatment right away may help keep these problems from becoming more serious. Your healthcare provider will check you for these problems during your treatment with TEVIMBRA. Your healthcare provider may treat you with corticosteroid or hormone replacement medicines, or may delay or completely stop treatment with TEVIMBRA if you have severe side effects.

Before receiving TEVIMBRA, tell your healthcare provider about all of your medical conditions, including if you have immune system problems; received an organ or tissue transplant; received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic); received radiation treatment to your chest area; or a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome; or are pregnant, plan to become pregnant, or are breastfeeding or plan to breastfeed. Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Can I take TEVIMBRA while pregnant or breastfeeding?

TEVIMBRA can harm your unborn baby. If you are a female who can become pregnant, your healthcare provider will do a pregnancy test before you start treatment with TEVIMBRA. Use effective birth control during treatment and for 4 months after your last dose. Do not breastfeed during treatment and for 4 months after your last dose. Talk to your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with TEVIMBRA.

The most common side effects of TEVIMBRA in combination with zanidatamab-hrii and fluoropyrimidine- and platinum-containing chemotherapy include diarrhea; nausea; decreased red blood cell count (anemia); decreased appetite; vomiting; decreased potassium in your blood; tiredness; rash; decreased white blood cell count; decreased platelet count; numbness, pain, tingling, or burning in your hands or feet; infusion reactions; and increase in certain liver blood tests.

These are not all the possible side effects of TEVIMBRA. Contact your healthcare provider for medical advice about side effects. You may report side effects to BeOne Medicines at 1-877-828-5596 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see additional Important Safety Information on the previous page and Medication Guide for TEVIMBRA.

TEVIMBRA®
(tislelizumab-jsgr)
Injection, for intravenous use 10 mg/mL

Visit TEVIMBRA.com to learn more

The information provided is meant for informational purposes only.
It is not meant to replace your physician's medical advice.

Actor portrayals.

BeOne

TEVIMBRA, BeOne Medicines, and myBeOneSupport are trademarks owned by BeOne Medicines I GmbH or its affiliates. All other trademarks are the property of their respective owners. © BeOne Medicines I GmbH, 2026 All Rights Reserved. 0526-TEV-PRC-212 08/26